

The Laplacian of a polynomial matrix of size $n \times n$ is symmetric

$$F = (x_1 + L_1^3, \dots, x_n + L_n^3) = (x_1, \dots, x_n) + L = x + L$$

$$L_i = \sum_{m=1}^n c_{im} x_m \quad c_{im} \in \mathbb{Z}$$

$$J_{ij} = \delta_{ij} + 3 L_i^2 \cdot L_{i,j} = \delta_{ij} + 3 L_i^2 c_{ij} = \delta_{ij} + 3 c_{ij} L_i^2$$

$$J_{ij} = J_{ji} \Rightarrow c_{ij} L_i^2 = c_{ji} L_j^2 \quad \forall i, j \in [n]$$

why should J be symmetric?? [Because, de Bruijnen, Esser, Druzkowski showed that it is enough to prove the conjecture with for symmetric Jacobian matrix.]

$$J = \begin{pmatrix} 1 + 3c_{11}L_1^2 & 3c_{12}L_1^2 & \dots & 3c_{1n}L_1^2 \\ 3c_{21}L_2^2 & 1 + 3c_{22}L_2^2 & \dots & 3c_{2n}L_2^2 \\ \vdots & \vdots & \ddots & \vdots \\ 3c_{n1}L_n^2 & 3c_{n2}L_n^2 & \dots & 1 + 3c_{nn}L_n^2 \end{pmatrix}$$

$$|J| = K \neq 1$$


$$\Rightarrow \text{there are lots of cancellations.} \quad c_{ij}L_i^2 = c_{ji}L_j^2$$

Idea: Graph of cancellations. (CANCELLATIONS)

~~why should the jacobian be symmetric??~~

$$J = \begin{pmatrix} 1 + 3c_{11}L_1^2 & 3c_{12}L_1^2 & \dots & 3c_{1n}L_1^2 \\ 3c_{12}L_2^2 & 1 + 3c_{22}L_2^2 & \dots & 3c_{2n}L_2^2 \\ \vdots & \vdots & \ddots & \vdots \\ 3c_{n1}L_1^2 & 3c_{n2}L_n^2 & \dots & 1 + 3c_{nn}L_n^2 \end{pmatrix}$$

rank $J = n \Rightarrow \exists \lambda_1, \dots, \lambda_n \in \mathbb{R}$ pairwise disjoint. $\forall x \in \mathbb{R}$
Let $\bar{v}_1, \dots, \bar{v}_n$ be the eigenvectors.



if $c_{ij} = 0$ then there is not such arrow. c_{ij} can be negative $\in \mathbb{Z}$.

Lemma:

~~$C_{ij} = 0 \Rightarrow C_{ji} = 0$~~

Proof:

$C_{ij} L_i^2 = C_{ji} L_j^2$

$\Rightarrow C_{ij} \left(\sum_{m=1}^n C_{im} x_m \right)^2 = C_{ji} \left(\sum_{m=1}^n C_{jm} x_m \right)^2$

$\Rightarrow C_{ij} = 0 \Rightarrow \left(C_{ji} = 0 \vee \sum_{m=1}^n C_{jm} x_m = 0 \right)$

$\Rightarrow C_{ji} = 0 \vee (C_{j1} = C_{j2} = \dots = C_{jn} = 0)$

$\Rightarrow C_{ji} = 0$

As there is symmetry between i and j , we also have the other implication

Lemma:

~~$C_{ij} = 0 \vee C_{ji} \neq 0$~~

~~$\frac{C_{ij}}{C_{ji}} = \frac{C_{jm_1} C_{jm_2}}{C_{im_1} C_{im_2}}$~~

~~$m_1, m_2 \in [n]$~~

~~they can be the same~~

~~$C_{ji} = \frac{C_{im_1} C_{im_2}}{C_{jm_1} C_{jm_2}} C_{ij}$~~

~~$m_1, m_2 \in [n]$~~

The following in particular

$C_{ij} (C_{im_1} C_{im_2}) = C_{ji} (C_{jm_1} C_{jm_2})$

$m_1, m_2 \in [n]$

Proof:

$C_{ij} L_i^2 = C_{ji} L_j^2$

they are allowed to be the same

Conway:

if $C_{ii} \neq 0 \vee i \in [n]$ then if and only if $C_{ij} = \pm C_{ji} \vee i \neq j \in [n]$

Proof:

~~$C_{ij} (C_{im_1} C_{im_2}) = C_{ji} (C_{jm_1} C_{jm_2})$~~

apply the previous lemma for $m_1 = i, m_2 = j$

$C_{ij} C_{ii} C_{ij} = C_{ji} C_{ji} C_{jj} \Rightarrow C_{ij}^2 = C_{ji}^2$

$C_{ij} C_{ii} C_{ij} = C_{ji} C_{ji} C_{jj} \Rightarrow C_{ij}^2 C_{ii} = C_{ji}^2 C_{jj}$

$\Rightarrow C_{ii} = C_{jj}$

QED

1: ~~Graph of F~~ L -graph of F : $G_L = (V, E)$
~~that~~ $V = \{x_1, \dots, x_n\}$.
 ~~$E = \{c_{ij}\}$~~ $e_{ij} = c_{ij}$ $E = \{e_{ij}\}$.

~~Corollary: If~~

~~Remark:~~

Corollary: If G_L has a constant λ then F is regular. $\lambda = 1$

Proof: If G_L has constant $\lambda \Rightarrow$
 $c_{ij} = 0 \iff i \neq j \Rightarrow \deg F = 1$
~~the~~ but the c_{ii} is free for degree 1-polynomial.

Remark: We need that all $c_{ii} \neq 0 \in \mathbb{C}[x]$.
 why? Because, otherwise, λ is not a constant. Hence $c_{ii} \in \mathbb{Z}^+ \forall i \in [n]$
 $\Rightarrow G_L$ has all the loops possible. if the coeff is false $\Rightarrow$ coeff max. with integer coeffs.

Task: Find interesting cases by looking at necessary condition on G_L for λ to be constant.
 $\rightarrow$ What is the determinant in terms of G_L ? Is there another graph G such that the Laplacian of $G = \lambda I$?
 (Not quite, as it would mean that G contains both c_{ij} and x_{ij} as ~~edges~~.)
 Maybe there is some graph product that gives us the coefficients of ~~x_{ij}~~ in $\det \lambda$.
 $x_{i_1} x_{i_2} \dots x_{i_z}$

Be careful, for this method is not suitable for fixed n .

$$b(a x_1 + b x_2)^2 = c(c x_1 + d x_2)^2$$

$$\begin{aligned} b a^2 &= c^3 \\ c d^2 &= b^3 \\ b^1 a &= c^1 d \end{aligned}$$

$$= 1 + 9a d (a x_1 + b x_2)^2 (c x_1 + d x_2)^2 + 3a (c x_1 + b x_2)^2 + 3d (c x_1 + d x_2)^2 - \cancel{3b^2} - 9b^2 (a x_1 + b x_2)^4$$

$$= 1 + (g_a d a^2 \cdot c^2 - g_b^2 a^4) x_1^4 + (g_a d b^2 \cdot d^2 - g_b^2 b^4) x_2^4 + \dots$$

Per MK.

~~It seems that
let $S = 1$ & abroad
S to
Then in really
known by people.
That in may be the
reason that ask
for this form 3
x 4 Li~~

Be careful for this mother!
is not suitable for fixed n!


It fits in the box, then my graph is selfless.

$$F_i = (x_1 + L_{i1}, \dots, x_n + L_{in})$$



$$J_{ij} = \delta_{ij} + 3 L_{ij}^2 \quad L_{ij} = \sum_{m=1}^n c_{ijm} x_m$$

$$\Rightarrow J_{ij} = \delta_{ij} + 3 \sum_{m=1}^n c_{ijm} x_m L_{ij}^2 = \delta_{ij} + 3 \sum_{m=1}^n c_{ijm} x_m^2 L_{ij}^2$$

$$J_{ij} = J_{ji} \Rightarrow c_{ij} L_{ij}^2 = c_{ji} L_{ji}^2$$



$$(a+b+c)(a+b+c) = a^2 + b^2 + c^2 + 2ab + 2ac + 2bc$$



$$\begin{cases} L_1 = c_{11}x_1 + c_{12}x_2 + c_{13}x_3 \\ L_2 = c_{21}x_1 + c_{22}x_2 + c_{23}x_3 \\ L_3 = c_{31}x_1 + c_{32}x_2 + c_{33}x_3 \end{cases}$$

$$c_{ij} (c_{i1}x_1 + c_{i2}x_2 + c_{i3}x_3)^2 = c_{ji} (c_{j1}x_1 + c_{j2}x_2 + c_{j3}x_3)^2$$

$$\text{If } c_{ij} = 0 \Rightarrow (c_{ji} = 0 \vee c_{j1} = c_{j2} = c_{j3} = 0)$$

$$c_{ij} \neq 0 : c_{ij} (c_{i1}^2 x_1^2 + c_{i2}^2 x_2^2 + c_{i3}^2 x_3^2 + 2c_{i1}c_{i2}x_1x_2 + 2c_{i1}c_{i3}x_1x_3 + 2c_{i2}c_{i3}x_2x_3) = c_{ji} (c_{j1}^2 x_1^2 + c_{j2}^2 x_2^2 + c_{j3}^2 x_3^2 + 2c_{j1}c_{j2}x_1x_2 + 2c_{j1}c_{j3}x_1x_3 + 2c_{j2}c_{j3}x_2x_3)$$

$$\Rightarrow \begin{cases} c_{ij} c_{i1}^2 = c_{ji} c_{j1}^2 \\ c_{ij} c_{i2}^2 = c_{ji} c_{j2}^2 \\ c_{ij} c_{i3}^2 = c_{ji} c_{j3}^2 \end{cases}$$

$$c_{ij} c_{i1} c_{i3} = c_{ji} c_{j1} c_{j3}$$

$$= \begin{pmatrix} g_1(T_1(x,y)) \\ g_2(T_2(x,y)) \end{pmatrix}$$

$$\begin{cases} T_{1x}(x,y)\dot{x} + T_{1y}(x,y)\dot{y} = g_1(T_1(x,y)) \end{cases}$$

$$\begin{cases} T_{2x}(x,y)\dot{x} + T_{2y}(x,y)\dot{y} = g_2(T_2(x,y)) \end{cases}$$

$$\begin{cases} T_{1x}(x,y)f_1(x,y) + T_{1y}(x,y)f_2(x,y) = g_1(T_1(x,y)) \end{cases}$$

$$\begin{cases} T_{2x}(x,y)f_1(x,y) + T_{2y}(x,y)f_2(x,y) = g_2(T_2(x,y)) \end{cases}$$

$$\begin{cases} z_x f_1 + z_y f_2 = g_1(z) \end{cases}$$

$$\begin{cases} w_x f_1 + w_y f_2 = g_2(w) \end{cases}$$



$$\dot{x}_4 = x_4^2$$

$$\dot{y} = x y$$

$$z = x^2$$

$$u = x y$$

$$\dot{z} = 2 x \dot{x} = \cancel{2 x^2} 2 x^3$$

$$\dot{u} = \dot{x} y + x \dot{y} = \cancel{2 x^2 y} \cancel{2 x^2 y}$$

$$\frac{d}{dt}(x y) = \frac{\partial(x y)}{\partial x} \dot{x} + \frac{\partial(x y)}{\partial y} \dot{y} = x^2 y + x^2 y = 2 x^2 y$$

$$\dot{z} = 2 x^3 = 2 x z = 2 z^{3/2}$$

$$\dot{u} = 2 x^2 y = 2 x u = 2 z^{1/2} u$$

$$x = \frac{1}{\sqrt{z}} u \quad T_1(x, y) = z$$

$$\dot{z} = \frac{\partial T_1}{\partial x} \dot{x} + \frac{\partial T_1}{\partial y} \dot{y} = \cancel{2 x^3}$$

$$\dot{u} = \frac{\partial T_2}{\partial x} \dot{x} + \frac{\partial T_2}{\partial y} \dot{y}$$

$$\begin{pmatrix} \dot{z} \\ \dot{u} \end{pmatrix} = \begin{pmatrix} T_{1x} & T_{1y} \\ T_{2x} & T_{2y} \end{pmatrix} \begin{pmatrix} \dot{x} \\ \dot{y} \end{pmatrix} = J_T \begin{pmatrix} \dot{x} \\ \dot{y} \end{pmatrix} = \begin{pmatrix} g_1(z) \\ g_2(u) \end{pmatrix}$$